

COBIA AQUACULTURE EXPANDING IN AMERICAS, CARIBBEAN

Summary:

A marine fish species with excellent characteristics for commercial aquaculture, cobia is receiving much attention in Latin American and Caribbean countries. Modern hatcheries are being built in the regions, and growout facilities are working to improve production efficiency. Continued success now should yield an expected boom in cobia culture in the near future.

Daniel Benetti, Ph.D.

Chairman, Division of Marine Affairs
and Policy
Associate Professor and Director
of Aquaculture
University of Miami
Rosenstiel School of Marine
and Atmospheric Science
4600 Rickenbacker Causeway
Miami, Florida 33149 USA
dbenetti@rsmas.miami.edu

Over the past decade, significant advances have been made in the culture of cobia, *Rachycentron canadum*. A marine fish species with excellent biological and market characteristics for commercial aquaculture, cobia has received much attention in many Latin American and Caribbean countries.

Research spearheaded by the University of Miami Experimental Hatchery in the last five years has progressed in the areas of broodstock procurement, management and spawning; larviculture; nutrition and sustainable growout technology. Much work remains to improve production efficiency, but countries throughout the Americas and the Caribbean – including Martinique, Puerto Rico, Dominican Republic, the United States, Mexico, Belize, Panama, Colombia, Ecuador and Brazil – are now producing cobia.



After successfully harvesting cobia from its Seastation cage system for five years, Snapperfarm celebrated the first commercial harvest from its Aquapod sea cage system in 2007. Cobia fingerlings initially grew at varied rates. Fingerling quality is of primary importance for growth performance. Photos by Brian O'Hanlon, Snapperfarm.

Albeit small compared to the numbers for other aquaculture species, global production of cobia is expected to exceed 25,000 mt in 2008 (mostly from Asia), with only about 500-1,000 mt produced in the United States, Caribbean and South America this year. The expanding industry is expected to boom in the next few years, when farmed cobia production in these regions could reach an estimated 2,000-3,000 mt within the next three years and 3,000-5,000 mt in five years.

New Hatcheries

Driven primarily by the anticipated cobia boom, new multimillion-dollar marine fish hatcheries are being built. In Brazil, four hatcheries in the states of São Paulo, Bahia, Espírito Santo and Pernambuco are working with cobia. In Belize, Marine Farms ASA – a company which already cultures cobia in Vietnam – is building a large-scale hatchery to produce cobia fingerling to stock a producing commercial cage farm.

In Panama, the large-scale commercial shrimp hatchery Farallón Aquaculture, which has facilities in several Latin American countries, is also constructing a large hatchery to produce cobia seedstock for the ex-

panding industry in that country. In Mexico, Ixoye Tropicales retrofitted an old shrimp hatchery to produce cobia fingerlings in 2007. A similar approach is being used by Ocean Farm in Ecuador, which adapted a large shrimp hatchery for the production of marine fish such as cobia, snapper and Pacific yellowtail.

The National Laboratory of Aquaculture, a state-of-the-art marine fish hatchery, just opened in Brazil. One of the most modern hatcheries on the continent, it is already in production utilizing technology transferred from the University of Miami through a memorandum of understanding with TWB S.A. and the Special Secretariat for Aquaculture and Fisheries, a partnership between the Brazilian private sector and government.

Similar cooperative agreements have been signed with Ecuador's Na-

**DRIVEN PRIMARILY
BY THE ANTICIPATED
COBIA BOOM, NEW
MULTIMILLION-
DOLLAR MARINE FISH
HATCHERIES ARE
BEING BUILT.**



The state-of-the-art hatchery at the National Laboratory of Aquaculture in Brazil incorporates a large-scale broodstock facility and associated infrastructure to support fingerling production for the country's growing cobia industry.

tional Center for Aquaculture and Marine Research (CENAIM-ESPOL) and Colombia's Center for Aquaculture Research (CENIACUA-ACUANAL) to further the development of marine fish aquaculture in these countries with technology transfer and training support. All these efforts will provide additional seedstock needed for industry expansion in the future.

Growout at Snapperfarm

Experiences gained at a commercial sea cage farm yielded varied and useful information on cobia growout. In 2006, Snapperfarm Inc. stocked cobia fingerlings from different hatcheries for growout in its cages off Culebra, Puerto Rico. Within a few weeks, each crop of fish from the various hatcheries was performing very differently. After six years growing cobia in its submerged cages, Snapperfarm found that growth rates for cobia can vary from 2 to 6 kg/year, depending on stocking density, temperature, feed quality and especially the quality of fingerlings.

One crop grew very well, while early signs of deformities were evident in the second crop and a third crop exhibited very slow growth. Genetics and nutritional fac-

tors may have affected the early development of the fish in the hatcheries. Also, Snapperfarm raised all three crops on different diets, in two different cage designs and at different densities – all of which could contribute to the varied performances early in the production cycle.

However, several months after stocking, all three crops were switched to a standard diet and the densities of two cages were matched, yet the performance still varied significantly from cage to cage. This led farm managers to believe that the quality of the hatchery-produced fingerlings is of paramount importance.

Despite these issues, Snapperfarm showed that properly planned, sited and operated open-ocean aquaculture operations can produce considerable amounts of high-quality seafood without causing significant environmental impacts. In fact, data from independent environmental assessments funded by the U.S. government being conducted by the University of Puerto Rico and University of Miami have shown no significant impact detected in oceanic waters surrounding the company's cages. This suggests that future expansion could occur in a sustainable manner.