

Simulations of the Arctic Basin with a finite-element sea-ice model

Olivier Lietaer^{1,2}, Vincent Legat¹, Thierry Fichefet²

¹ Centre for Systems Engineering and Applied Mechanics (CESAME),
Université catholique de Louvain, Louvain-la-Neuve, Belgium.

² G. Lemaitre Institute of Astronomy and Geophysics (ASTR),
Université catholique de Louvain, Louvain-la-Neuve, Belgium.

September 29, 2006

Oral presentation

Abstract

We are currently developing a finite element model to simulate large-scale sea-ice dynamics in an Eulerian description. Our model has representations of both dynamic and thermodynamic sea-ice processes, and computes the ice pack velocity pattern and its thickness and concentration distribution. Sea-ice is assumed to behave as a two-dimensional viscous-plastic continuum in dynamical interaction with atmosphere and ocean. The vertical growth/melt rate of the ice is computed by the zero-layer model of *Semnter* (1976). The thermodynamic processes include a parameterization of the different heat fluxes interacting with the ice. A simulation of the Arctic ice pack is carried out on a realistic mesh by using NCEP/NCAR reanalysis of the atmospheric fields to drive our model. We compare the simulated ice velocity, thickness and concentration patterns with those derived from satellite and buoy measurements. Finally, a study of the grid resolutions influence on the results is presented.

References

Semnter, A. J. (1976). A model for the thermodynamic growth of sea ice in numerical investigations of climate. *Journal of Physical Oceanography*, 6, 379-389.